

Part 1. Biotechnology and genetics

UDC 619:578.828:578.2'21:636.22/.28

STUDYING OF PHYLOGENETIC RELATIONSHIPS OF LEUKEMIA VIRUS WITH OTHER RETROVIRUSES IN CATTLE

Limanskaya O. Yu., Gema I. A., Gorbatenko S. K., Gerilovych A. P.

National Scientific Center 'Institute of Experimental and Clinical Veterinary Medicine',
Kharkiv, Ukraine, e-mail: antger2011@gmail.com

Summary. Bovine leukemia virus (BLV) is one of the retroviruses, which genetically, structurally and functionally related to viruses of human T-cell leukemia. BLV is a very convenient model for studying the pathogenesis of human leukemia. Genomes of retroviruses have high variability levels due to lack of a mechanism of correct errors that occur when copying matrix during replication, and possible genetic recombination. In this regard, the study of the genetic variability of the virus is one of the major objectively for biological monitoring. At this time, molecular genetic analysis (polymerase chain reaction (PCR)) is a necessary part of phylogenetic research. The aim of this work was to study the variability of the bovine leukemia virus, to establish phylogenetic relationships between isolates sequenced bovine leukemia virus, which circulates in farms of different regions in Ukraine, with other animals retroviruses. The sampling of clinical material from cattle farms was conducted in different geographical regions in Ukraine and extracted proviral BLV DNA. Totally 831 samples of peripheral blood were collected and tested from cattle farms in Kharkiv region, 10 samples — Kirovohrad region; 10 samples — Donetsk region; 41 samples — Crimea, Simferopol region; 10 — samples of Poltava region. Sequenced fragments of env gene of bovine leukemia virus proviral DNA, circulating in different geographical regions in Ukraine were analyzed. Established isolates of bovine leukemia virus, circulating in Ukraine, belonging to the Euro-Asian subtype. Proved genetic affinity of leukemia virus and bovine syncytial virus, Jembrana disease virus and bovine immunodeficiency virus.

Keywords: DNA, phylogeographic relationships, polymerase chain reaction, sequencing, retroviruses, virus bovine leukemia.

Introduction. Bovine leukemia virus (BLV) is the one of the retroviruses representatives. Humans T-cell leukemia viruses, HTLV-1 and HTLV-2, genetically, structurally and functionally related to bovine leukemia virus, and the development of diseases caused by these viruses is similar (Willems et al., 1993; Dube et al., 1997). That is why BLV is a very suitable model for studying the pathogenesis of human leukemia. An important aspect of these studies is the problem of bovine leukemia virus genetic variability.

It is known that viruses, which genome is represented by RNA, characterized by high-speed variation of nucleotide sequence and associated with it significant lability structure of genetic material (Steinhauer and Holland, 1987; Parvin et al., 1986; Steinhauer et al., 1993; Darlix and Spahr, 1983; Katz and Skalka, 1990; Meyerhans et al., 1989; Manini, De Palma, and Mutti, 2007). Genomes of retroviruses, like other RNA-containing viruses, have high level of variability due to lack of mechanism for correcting errors arising until copying the matrix during replication and potential genetic recombination. Nucleotide modifications can

lead to changes an amino acid consist of synthesized proteins (Katz and Skalka, 1990). In this regard, the study of genetic variability of infectious agents is the one of the main objectives of biological monitoring, that goal is the explanation of the phenomenon.

At this time, the necessary part of phylogenetic study is molecular genetic analysis (such as polymerase chain reaction (PCR) (Licursi et al., 2003; Giammarioli et al., 2008). The method of sequencing allows performing of the existence of point and tandem mutations (Milos, 2009). Effectiveness and objectivity of molecular phylogenetic studies depend on many factors, such as: insufficient set of experimental data, errors in sequencing or sequence alignment, convergent evolution (i.e. formation of a complex of similar features in representatives from unrelated groups), horizontal gene transfer, etc. (Wendel and Doyle, 1998). In addition, different fragments of the genome provide unequal information during the molecular phylogenetic studies, because the result is more determined by the correct choice of gene or combination of genes in the sequence. Sometimes complementary DNA (cDNA), that includes

coding sequences of structural genes, used for sequencing (Caraguel et al., 2009; Chang et al., 2009).

The aim of this work was to study the variability of the bovine leukemia virus, to establish phylogenetic relationships of sequenced isolates bovine leukemia virus that circulates in farms of different regions in Ukraine with other animals retroviruses.

Materials and methods. The total DNA was extracted from peripheral blood using a commercial kit 'DNA Sorbo-B' (Russia). Detection of proviral DNA was performed by PCR using basic kits Gene Pak™ (Russia) and a pair of primers III-BLV F / R developed in 2008. The length of the amplicon is 440 bp.

Sequencing of proviral DNA of *env* gene fragments was performed on an automatic DNA-analyzer ABI PRISM 311D using the technology of ABI (Applied Biosystems, USA).

Computer analysis of the primary structure of bovine leukemia virus isolates proviral DNA fragments, multiple alignment of proviral DNA sequenced of major genes of retroviruses was carried out using programs Bioedit (ClustalW modules and Neighbor), version 7.0.0, and Oligo Explorer, version 1.1.0. To construct phylogenetic trees used the program MEGA, version 4.1., and to view them — TreeView, version 1.6.6.

Results. The sampling of clinical material of cattle was performed in farms of different geographical regions in Ukraine and proviral DNA of BLV was extracted. There were analyzed 831 samples of peripheral blood of cattle from Kharkiv region farms, 10 samples — Kirovohrad region; 10 samples — Donetsk region; 41 samples — Crimea, Simferopol region; 10 samples — Poltava region.

To set the primary structure of fragments of BLV genomic material by sequencing, the samples of proviral DNA were transferred to the National Veterinary Research Institute (Pulawy, Poland).

Multiple alignment and comparison of BLV *env* gene sequences circulating in the Kharkiv region and other geographical regions was created to determine the possible divergence and the genetic variability of bovine leukemia virus. For this purpose from the international database GenBank we have selected three fully sequenced BLV *env* gene sequences from Belgium (AF503581), USA (AY078387) and Brazil (AF399704) with length 1548 bp, and partially sequenced fragment with length 960 bp from Poland (AF111171). Fragment of the multiple alignment of selected sequences demonstrated in Fig. 1 (nucleotides that are not the same in this position with other nucleotides of selected sequences are underlined).

	425	435	445	455	465	475
AF111171	GGCGATCAAA	GGGTCTCTTT	ATGTCAATCA	TCAGATTITA	TTCTGCAATC	TCAAAACATG
AF399704	GGCGATCAAA	GGATCTCTTT	ATGTCAATCA	TCAGATTITA	TTCTGCAATC	TCAAAACATG
AF503581	GGCGATCAAA	GGGTCTCTTT	ATGTCAATCA	TCAGATTITA	TTCTGCAATC	TCAAAACATG
AY078387	GGCGATCAAA	GGATCTCTTT	ATGTCAATCA	TCAGATTITA	TTCTGCAATC	TCAAAACATG
Kharkiv	GGCGATCA <u>AG</u>	GGATCTCTTT	ATGTCAATCA	TCAGATTITA	TTCTGCAATC	TCAAAACATG
	485	495	505	515	525	535
AF111171	TCATGGAAAT	TTCACTCTAA	CCTGGGAAAT	ATGGGGATAT	GATCCCTGTA	TCACCTTTTC
AF399704	TCATGGAAAT	TTCACTTTAA	CCTGGGAAAT	ATGGGGATAT	GATCCCTGTA	TCACCTTTTC
AF503581	TCATGGAAAT	TTCACTCTAA	CCTGGGAAAT	ATGGGGATAT	GATCCCTGTA	TCACCTTTTC
AY078387	TCATGGAAAT	TTCACTCTAA	CCTGGGAAAT	ATGGGGATAT	GATCCCTGTA	TCACCTTTTC
Kharkiv	TCATGGAAAT	TTCACTCTAA	CCTGGGAAAT	ATGGGGATAT	GATCCCTGTA	TCACCTTTTC
	545	555	565	575	585	595
AF111171	TTTACATAAA	ATCCCTGATC	CCCTCAACC	CGACTTCCTT	CAGCTGAACA	GTGACTGGGT
AF399704	TTTACATAAG	ATCCCTGATC	CCCTCAACC	CGACTTCCTT	CAGCTGAACA	GTGACTGGGT
AF503581	TTTACATAAA	ATCCCTGATC	CCCTCAACC	CGACTTCCTT	CAGCTGAACA	GTGACTGGGT
AY078387	TTTACATAAG	ATCCCTGATC	CCCTCAACC	CGACTTCCTT	CAGCTGAACA	GTGACTGGGT
Kharkiv	TTTACATAAG	ATCCCTGATC	CCCTCAACC	CGACTTCCTT	CAGCTGAACA	GTGACTGGGT
	605	615	625	635	645	655
AF111171	TCCCTCTGTC	AGATCATGGG	CCCTGCTTTT	AAATCAACCG	GCAAGGGGCT	TCCAGACTG
AF399704	TCCCTCTGTC	AGATCATGGG	CCCTGCTTTT	AAATCAACCG	GCAAGGGGCT	TCCAGACTG
AF503581	TCCCTCTGTC	AGATCATGGG	CCCTGCTTTT	AAATCAACCG	GCAAGGGGCT	TCCAGACTG
AY078387	TCCCTCTGTC	AGATCATGGG	CCCTGCTTTT	AAATCAACCG	GCAAGGGGCT	TCCAGACTG
Kharkiv	TCCCTCTGTC	AGATCATGGG	CCCTGCTTTT	AAATCAACCG	GCAAGGGGCT	TCCAGACTG
	665	675	685	695	705	715
AF111171	TGCTATATGT	TGGGAACCTT	CCCTCCCTG	GGCTCCCGAA	ATATTAGTAT	ATAACAAAAC
AF399704	TGCTATATGT	TGGGAACCTT	CCCTCCCTG	GGCTCCCGAA	ATATTAGTAT	ATAACAAAAC
AF503581	TGCTATATGT	TGGGAACCTT	CCCTCCCTG	GGCTCCCGAA	ATATTAGTAT	ATAACAAAAC
AY078387	TGCTATATGT	TGGGAACCTT	CCCTCCCTG	GGCTCCCGAA	ATATTAGTAT	ATAACAAAAC
Kharkiv	TGCTATATGT	TGGGAACCTT	CCCTCCCTG	GGCTCCCGAA	ATATTAGTAT	ATAACAAAAC
	725	735	745	755	765	775
AF111171	CATCTCCAGC	TCTGGACCGG	GTCTCGCCCT	CCCGAGCGCC	CAAAATCTTCT	GGGTCAACAC
AF399704	CATCTCCAGC	TCTGGACCGG	GTCTCGCCCT	CCCGAGCGCC	CAAAATCTTCT	GGGTCAACAC
AF503581	CATCTCCAGC	TCTGGACCGG	GTCTCGCCCT	CCCGAGCGCC	CAAAATCTTCT	GGGTCAACAC
AY078387	CATCTCCAGC	TCTGGACCGG	GTCTCGCCCT	CCCGAGCGCC	CAAAATCTTCT	GGGTCAACAC
Kharkiv	CATCTCCAGC	TCTGGACCGG	GTCTCGCCCT	CCCGAGCGCC	CAAAATCTTCT	GGGTCAACAC

Figure 1. Fragment of multiple sequence alignment of *env* gene of bovine leukemia virus circulating in different geographical regions

The same value of divergence of the BLV *env* gene, of agent circulating in the Kharkiv region, characterized proviral DNA sequences from the agents circulating in western Europe (Table 1). The smallest divergence levels were observed for the *env* gene of European isolates of bovine leukemia virus. In general, the results supported that the BLV *env* gene is highly conserved, and its primary structure does not change depending on the habitat of the causative agent.

Table 1 — The degree of similarity and divergence of the *env* gene of BLV circulating in different geographic regions, calculated relative to the corresponding sequence from isolate fragments of Kharkiv region

	AF 111171 (Poland)	AF 399704 (Brazil)	AF 503581 (Belgium)	AY 078387 (USA)
The number of nucleotides that do not coincide	14	15	14	19
Divergence, %	1.8	2.0	1.8	2.5
The degree of similarity, %	98.2	98	98.2	97.5

The phylogenetic tree, that illustrated proximity of BLV isolates circulating in Ukraine to isolates of European and Asian subgroups was constructed to establish phylogenetic relations between isolates of bovine leukemia virus circulating in Ukraine, and their phylogenetic relationships with isolates from other regions of the world (Europe, Asia, North and South America), (1 in the Fig. 2). BLV isolates, that proviral DNA was extracted from peripheral blood of animals from farms in Rivno, Poltava and Kharkiv regions, are closer to the European subgroup (Austria isolate). Bovine leukemia virus, circulating in the farms of Crimea, is closer to Asian subgroup (Zanjan, Tehran isolates). Isolates from the America form a separate, American, subgroup (2 in the Fig. 2).

Thus, the results of phylogenetic studies can be used to identify and study possible subgroups (or genotypes), to create the basis for the genes search, that determine the high biological activity of viruses.

To study the phylogenetic relationships of bovine leukemia virus was created databases of sequenced gene sequences and their fragments, isolated

in different geographical regions and represented in international databases GenBank and EMBL: the bovine immunodeficiency virus, the Jembrana disease virus — lentivirus, that causes severe acute disease of cattle characterized by lymphopenia and lymphadenopathy; syncytial virus; bovine leukemia virus.




Figure 2. Dendrogram, based on *env* gene fragment of proviral DNA of bovine leukemia virus isolates from different geographical regions

Phylogenetic analysis, based on retroviruses *env* gene sequences (Fig. 3) and completely sequenced proviral DNA sequences (Fig. 4), showed firstly that isolates of Jembrana disease viruses, immunodeficiency and syncytial viruses form separate branches; secondly, membership of bovine leukemia virus isolates to one cluster; thirdly, the evolutionary closeness of the leukemia virus and bovine syncytial virus.




Figure 3. Dendrogram, based on *env* gene fragment of proviral DNA of bovine retroviruses isolates




Figure 4. Dendrogram, based on complete sequences of proviral DNA of bovine retroviruses isolates

Conclusions. The BLV env gene is highly conserved, its primary structure has significant changes depending on the habitat of the causative agent of bovine enzootic leukosis. The greatest degree of similarity observed for env gene sequences of European BLV isolates. It was shown, that bovine leukemia virus isolates, circulating in farms of different geographic regions in Ukraine, are closer to the Euro-Asian subgroup. Based on phylogenetic analysis, proved the genetic proximity of leukemia virus and bovine syncytial virus, immunodeficiency virus and Jembrana diseases virus.

References

- Caraguel, C., Stryhn, H., Gagné, N., Dohoo, I. and Hammell, L. (2009) 'Traditional descriptive analysis and novel visual representation of diagnostic repeatability and reproducibility: Application to an infectious salmon anaemia virus RT-PCR assay', *Preventive Veterinary Medicine*, 92(1–2), pp. 9–19. doi: 10.1016/j.prevetmed.2009.07.011.
- Chang, Z., Jin, M., Liu, N., Xie, H., Cui, S., Zhang, Q. and Duan, Z. (2009) 'Analysis of epidemiologic feature and genetic sequence of Sapovirus in China', *Chinese Journal of Virology*, 25(2), pp. 113–116. doi: 10.13242/j.cnki.bingduxuebao.001979.
- Darlix, J.-L. and Spahr, P.-F. (1983) 'High spontaneous mutation rate of Rous sarcoma virus demonstrated by direct sequencing of the RNA genome', *Nucleic Acids Research*, 11(17), pp. 5953–5967. doi: 10.1093/nar/11.17.5953.
- Dube, S., Bachman, S., Poiesz, B. J., Ferrer, J. F., Esteban, E., Choi, D., Love, J. and Spicer, T. (1997) 'Degenerate and specific PCR assays for the detection of bovine leukaemia virus and primate T cell leukaemia/lymphoma virus pol DNA and RNA: Phylogenetic comparisons of amplified sequences from cattle and primates from around the world', *Journal of General Virology*, 78(6), pp. 1389–1398. doi: 10.1099/0022-1317-78-6-1389.
- Giammarioli, M., Pellegrini, C., Casciari, C., Rossi, E. and De Mario, G. M. (2008) 'Genetic diversity of bovine viral diarrhoea virus 1: Italian isolates clustered in at least seven subgenotypes', *Journal of Veterinary Diagnostic Investigation*, 20(6), pp. 783–788. doi: 10.1177/104063870802000611.
- Katz, R. A. and Skalka, A. M. (1990) 'Generation of diversity in retroviruses', *Annual Review of Genetics*, 24(1), pp. 409–443. doi: 10.1146/annurev.ge.24.120190.002205.
- Licursi, M., Inoshima, Y., Wu, D., Yokoyama, T., González, E. T. and Sentsui, H. (2003) 'Provirus variants of bovine leukemia virus in naturally infected cattle from Argentina and Japan', *Veterinary Microbiology*, 96(1), pp. 17–23. doi: 10.1016/s0378-1135(03)00202-5.
- Manini, P., De Palma, G. and Mutti, A. (2007) 'Exposure assessment at the workplace: Implications of biological variability', *Toxicology Letters*, 168(3), pp. 210–218. doi: 10.1016/j.toxlet.2006.09.014.
- Meyerhans, A., Cheynier, R., Albert, J., Seth, M., Kwok, S., Sninsky, J., Morfeldt-Månson, L., Asjö, B. and Wain-Hobson, S. (1989) 'Temporal fluctuations in HIV quasiespecies in vivo are not reflected by sequential HIV isolations', *Cell*, 58(5), pp. 901–910. doi: 10.1016/0092-8674(89)90942-2.
- Milos, P. M. (2009) 'Emergence of single-molecule sequencing and potential for molecular diagnostic applications', *Expert Review of Molecular Diagnostics*, 9(7), pp. 659–666. doi: 10.1586/erm.09.50.
- Parvin, J. D., Moscona, A., Pan, W. T., Leider, J. M. and Palese, P. (1986) 'Measurement of the mutation rates of animal viruses: Influenza A virus and poliovirus type 1', *Journal of Virology*, 59(2), pp. 377–383. Available at: <http://jvi.asm.org/content/59/2/377.full.pdf>.
- Steinhauer, D. A. and Holland, J. J. (1987) 'Rapid evolution of RNA viruses', *Annual Review of Microbiology*, 41(1), pp. 409–431. doi: 10.1146/annurev.mi.41.100187.002205.
- Steinhauer, D. A., Torre, J. C. de la, Meier, E. and Holland, J. J. (1989) 'Extreme heterogeneity in populations of vesicular stomatitis virus', *Journal of Virology*, 63(5), pp. 2072–2080. Available at: <http://jvi.asm.org/content/63/5/2072.full.pdf>.
- Wendel, J. F. and Doyle, J. J. (1998) 'Phylogenetic incongruence: Window into genome history and molecular evolution. In: Soltis, D. E., Soltis, P. S. and Doyle, J. J. *Molecular Systematics of Plants II*. Boston: Kluwer Academic Publishing. doi: 10.1007/978-1-4615-5419-6_10.
- Willems, L., Thienpont, E., Kerkhofs, P., Burny, A., Mammerickx, M. and Kettmann, R. (1993) 'Bovine leukemia virus, an animal model for the study of intrastrain variability', *Journal of Virology*, 67(2), pp. 1086–1089. Available at: jvi.asm.org/content/67/2/1086.full.pdf.