

Part 2. Biotechnology

UDC 619:616.98-078:578.82/.83[PCV-3]:577.2.08:602.6:636.4

DOI 10.36016/JVMBBS-2021-7-3-3

DEVELOPMENT OF RECOMBINANT POSITIVE CONTROL FOR DETECTION OF PORCINE CIRCOVIRUS TYPE 3 BY POLYMERASE CHAIN REACTION

Rudova N. G., Lymanska O. Yu., Bolotin V. I., Stegnyy B. T., Solodiankin O. S., Gerilovych A. P.

National Scientific Center 'Institute of Experimental and Clinical Veterinary Medicine', Kharkiv, Ukraine, e-mail: rudovanatawa@ukr.net

Summary. This work aimed to obtain positive control using recombinant DNA technology for detection by PCR of a new poorly studied pathogen — porcine circovirus type 3. Recombinant positive control was designed using Clone Manager Basic. As a vector in the creation of recombinant control we used plasmid pTZ57R/T, as an insert — a fragment of the gene *rep* PCV-3 with the length of 418 nucleotide pairs, obtained by classical PCR. Transformation of competent cells of *E. coli* strain DH5a was carried out by chemical poration, followed by plating on LB-medium with the addition of ampicillin at a final concentration of 100 µg/ml. The selection of *E. coli* cell colonies was performed by the marker of antibiotic resistance to ampicillin. The presence of a specific insert was checked by PCR with electrophoretic visualization of the results. The developed recombinant positive control can be used for the monitoring of biological samples from pigs for the presence of genetic material PCV-3 using molecular technologies

Keywords: PCV-3, PCR, plasmid pTZ57R/T, gene *rep*, *E. coli* strain DH5a

Introduction. One of the branches of animal husbandry that is actively developing all over the world, including Ukraine, is pig breeding. The level of development of this industry and the quality of the products obtained depend on the welfare of pig breeding concerning infectious animal diseases. One of the most common diseases among pigs today is circovirus infection, which causes significant economic damage to the industry and is characterized by severe immunosuppression and multiorgan pathology. Losses from PCV infection are estimated by the death of piglets at the stages of rearing and fattening; failure to gain weight of piglets; losses due to pathology of reproduction; ineffectiveness of antibiotic therapy in the presence of bacterial coinfections.

The causative agent of PCV disease is mainly porcine circovirus type 2, which belongs to the family Circoviridae, is characterized by pronounced pathogenicity and is the cause of many syndromes and diseases of pigs. In 2016, it was reported about a new type of porcine circovirus — PCV-3 in sows with clinical signs of dermatitis and nephropathy, as well as with reproductive problems (Palinski et al., 2017).

Currently known data indicate the prevalence of PCV-3 among pigs and wild boar populations in many countries in different geographical regions of the world (Bera et al., 2020; Hayashi et al., 2018; Serena et al., 2021; Saraiva et al., 2019; Saporiti et al., 2020b; Yuzhakov et al., 2018; Franzo et al., 2018; Souza et al., 2021).

PCV-3 is associated with cardiac and multisystem inflammations, dermatitis and nephropathy syndrome, abortion and reproductive disorders in pigs, respiratory diseases; it is able to cause intrauterine infections in the absence of obvious reproductive disorders (Jiang et al.,

2019; Ouyang et al., 2019; Phan et al., 2016; Saporiti et al., 2020c).

At present, for the detection of PCV-3 and rapid assessment of the epizootic situation, methods based on amplification technologies, including the use of different formats of polymerase chain reaction (PCR), is mainly used (Ji et al., 2019; Liu et al., 2019; Wang et al., 2019; Kim et al., 2020; Yuan et al., 2020; Zheng et al., 2020).

An important component that ensures the high quality of molecular genetic tests based on PCR and the level of reliability of the results obtained regarding the detection of PCV-3 is a positive control, which often uses material containing the virus. The use of such control is caused by the need to periodically obtain the appropriate material with a limited shelf life.

Therefore, the **aim of this work** was to construct a recombinant positive control containing an insert of gene *rep* PCV-3 with a length of 418 bp.

Materials and methods. Virtual design of recombinant positive control was performed using Clone Manager Basic v. 9 (Sci Ed Software, USA).

Plasmid vector pTZ57R/T, which is part of the commercial kit 'InsTAclone PCR Cloning Kit' (Fermentas, Latvia), was used to create *in vitro* plasmid control.

A fragment of the *rep* PCV-3 gene with a length of 418 bp was used as an insert. It was obtained by classical PCR using the commercial 'Maxima Hot Start Green PCR Master Mix' (Thermo Scientific, Lithuania) and the PCV-3 primer system F (5'-TTGTGGTGCTACGAGTG TCC-3'); PCV-3 R (5'-CGTCTCCGTCAGAATCCGAG-3') (Saporiti et al., 2020a).

Integration of the plasmid molecule into the culture of competent cells of *E. coli* strain DH5a was performed

by chemical poration followed by plating on LB medium (Sigma-Aldrich, USA) with the addition of ampicillin at a final concentration of 100 µg/ml.

A commercial kit 'Plasmid Miniprep Kit' (GeneJET, Lithuania) was used to extract the plasmid DNA.

Measurement of DNA concentration and evaluation of DNA quality was performed using a NanoDrop spectrophotometer (DeNovix, USA).

Electrophoretic analysis of amplification products was performed by horizontal electrophoresis (horizontal electrophoresis chamber Biorad (USA) in 1.5% agarose gel.

For electrophoretic analysis we used agarose produced by Biozym (Germany), ethidium bromide manufactured by Sigma-Aldrich (USA), molecular weight markers manufactured by Invitrogen (USA), Promega (USA) and Fermentas Gene ruler (Latvia) with a discreteness of 100 bp.

Results. In order to obtain a recombinant positive control sample for the detection of PCV-3 genetic material by PCR, we constructed a virtual model of a vector molecule based on plasmid pTZ57R/T with an embedded fragment of the *rep* PCV-3 gene with a length of 418 bp. The total length of the theoretically modeled plasmid molecule was 3,304 bp (Fig. 1).




Figure 1. Scheme of recombinant plasmid pTZ57R/T_PCV-3

To create a plasmid control *in vitro* in the first stage of our work, a fragment of the *rep* PCV-3 gene with a length of 418 bp was developed. To do this, we used a DNA sample obtained from pig liver homogenate, which we had previously described as positive for the presence of PCV-3 genetic material. The analysis by horizontal gel electrophoresis confirmed the presence of an amplicon of the required length — 418 bp.

The resulting amplicon was purified and ligated to the plasmid vector pTZ57R/T, which transformed competent *E. coli* DN5a cells.

Because this vector contained an ampicillin resistance gene, it was a marker of selective traits during subsequent cloning of strain DN5a in *E. coli* culture.

Therefore, after the transformation, six white single colonies of *E. coli* with signs of acquired resistance to ampicillin were selected (Fig. 2).




Figure 2. Selection of colonies of *E. coli* cells by the marker of antibiotic resistance to ampicillin

PCR screening of selected colonies showed the presence of a specific insert with the length of 418 bp in each of them (Fig. 3).

agarose gel of the obtained samples testified to the successful transformation of *E. coli* cells by the recombinant plasmid constructed by us (Fig. 4).




Figure 3. Visualization of the result of PCR screening of selected colonies by electrophoretic analysis in 1.5% agarose gel (M — molecular weight marker, NC — negative control, PC — positive control, 1–6 — samples positive for the presence of PCV-3 DNA)

Three colonies of *E. coli* were selected for cultivation in liquid nutrient medium, after which the resulting bacterial biomass was used for extraction of plasmid DNA. The presence of fragments about 3 thousand bp in length when performing electrophoretic analysis in 1.5%




Figure 4. Visualization of the result of electrophoretic analysis of purified plasmid DNA (M — molecular weight marker; 1–3 — positive for the presence of plasmid DNA samples)

The concentration of DNA in the first sample was 106.15 ng/μl, in the second — 132.39 ng/μl, in the third — 83.08 ng/μl (Fig. 5).




Figure 5. Results of plasmid DNA concentration measurement

The sample with the highest concentration of plasmid DNA was selected for our further studies.

Thus, we have developed a recombinant sample that contains a fragment with a length of 418 bp porcine circovirus type 3 *rep* gene. It can be used as a positive control in the detection of PCV-3 by PCR.

Discussion. Issues related to the selection and design of adequate positive controls for different PCR formats have always received considerable attention, primarily due to the potential risk to biosafety during the study (Chan, Jiang and Tan, 2016; Caasi et al., 2013; Chen et al., 2006; Lion, 2001).

Considerable factors in ensuring the high accuracy and efficiency of molecular genetic analysis based on PCR are, in particular, the conditions and form of storage of biological material used as a positive control.

At present, the use of positive control in the form of a recombinant plasmid containing a fragment of certain genomic DNA has become widespread in the detection of infectious diseases in farm animals, poultry (Yao et al., 2019; Das et al., 2017) and humans (Camacho et al., 2016), food pathogens (Gokduman et al., 2016), food allergens (Miyazaki et al., 2019), determination of GMOs (Taverniers, Van Bockstaele and De Loose, 2004) due to the long shelf life of such a structure, high copy capacity, the possibility of recovery (Matange, Tuck and Keung, 2021).

An equally important advantage of recombinant positive controls is that they do not require the permits provided by the Cartagena Protocol on Biosafety to the Convention on Biological Diversity, which was signed by Ukraine in 1992. Therefore, the recombinant positive control developed by us can be used to monitor biological samples obtained from pigs for the presence of PCV-3 genetic material using molecular technologies.

Conclusions. Thus, we obtained an ampicillin-resistant clone of *E. coli* DN5 α , transformed with the constructed plasmid pTZ57R/T_PCV-3 with insertion of the *rep* gene with a length of 418 bp, which can be used as a positive control sample for the detection of the PCV-3 genetic material by PCR method.

References

- Bera, B. C., Choudhary, M., Anand, T., Virmani, N., Sundaram, K., Choudhary, B. and Tripathi, B. N. (2020) 'Detection and genetic characterization of porcine circovirus 3 (PCV3) in pigs in India', *Transboundary and Emerging Diseases*, 67(3), pp. 1062–1067. doi: 10.1111/tbed.13463.
- Caasi, D. R. J., Arif, M., Payton, M., Melcher, U., Winder, L. and Ochoa-Corona, F. M. (2013) 'A multi-target, non-infectious and clonable artificial positive control for routine PCR-based assays', *Journal of Microbiological Methods*, 95(2), pp. 229–234. doi: 10.1016/j.mimet.2013.08.017.
- Camacho, D., Reyes, J., Franco, L., Comach, G. and Ferrer, E. (2016) 'Cloning alphavirus and flavivirus sequences for use as positive controls in molecular diagnostics' [Clonación de secuencias de alfavirus y flavivirus para uso como controles positivos en el diagnóstico molecular], *Revista Peruana de Medicina Experimental y Salud Pública*, 33(2), pp. 269–273. doi: 10.17843/rpmpesp.2016.332.2101. [in Spanish].
- Chan, M., Jiang, B. and Tan, T.-Y. (2016) 'Using pooled recombinant plasmids as control materials for diagnostic real-time PCR', *Clinical Laboratory*, 62(10), p. 1893–1901. doi: 10.7754/Clin.Lab.2016.160114.
- Chen, J.-M., Guo, L.-X., Sun, C.-Y., Sun, Y.-X., Chen, J.-W., Li, L. and Wang, Z.-L. (2006) 'A stable and differentiable RNA positive control for reverse transcription-polymerase chain reaction', *Biotechnology Letters*, 28(22), pp. 1787–1792. doi: 10.1007/s10529-006-9161-0.
- Das, A., Ward, G., Lowe, A., Xu, L., Moran, K., Renshaw, R., Dubovi, E., Reising, M. and Jia, W. (2017) 'Development and validation of a highly sensitive real-time PCR assay for rapid detection of parvoviruses', *Journal of Veterinary Diagnostic Investigation*, 29(4), pp. 499–507. doi: 10.1177/1040638716680676.
- Franzo, G., Tucciarone, C. M., Drigo, M., Cecchinato, M., Martini, M., Mondin, A. and Menandro, M. L. (2018) 'First report of wild boar susceptibility to porcine circovirus type 3: High prevalence in the Colli Euganei Regional Park (Italy) in the absence of clinical signs', *Transboundary and Emerging Diseases*, 65(4), pp. 957–962. doi: 10.1111/tbed.12905.
- Gokduman, K., Dilek Avsaroglu, M., Cakiris, A., Ustek, D. and Candan Gurakan, G. (2016) 'Recombinant plasmid-based quantitative real-time PCR analysis of *Salmonella enterica* serotypes and its application to milk samples', *Journal of Microbiological Methods*, 122, pp. 50–58. doi: 10.1016/j.mimet.2016.01.008.
- Hayashi, S., Ohshima, Y., Furuya, Y., Nagao, A., Oroku, K., Tsutsumi, N., Sasakawa, C. and Sato, T. (2018) 'First detection of porcine circovirus type 3 in Japan', *Journal of Veterinary Medical Science*, 80(9), pp. 1468–1472. doi: 10.1292/jvms.18-0079.
- Ji, J., Xu, X., Wang, X., Zuo, K., Li, Z., Leng, C., Kan, Y., Yao, L. and Bi, Y. (2019) 'Novel polymerase spiral reaction assay for the visible molecular detection of porcine circovirus type 3', *BMC Veterinary Research*, 15(1), p. 322. doi: 10.1186/s12917-019-2072-9.
- Jiang, H., Wang, D., Wang, J., Zhu, S., She, R., Ren, X., Tian, J., Quan, R., Hou, L., Li, Z., Chu, J., Guo, Y., Xi, Y., Song, H., Yuan, F., Wei, L. and Liu, J. (2019) 'Induction of Porcine dermatitis and nephropathy syndrome in piglets by infection with porcine circovirus type 3', *Journal of Virology*, 93(4), p. e02045-18. doi: 10.1128/JVI.02045-18.
- Kim, H., Lim, D., Chae, H., Park, J., Kim, S., Lee, K., Lee, C., Lyoo, Y. S. and Park, C. (2020) 'Advanced target-specific probe-based real-time loop-mediated isothermal amplification assay for the rapid and specific detection of porcine circovirus 3', *Transboundary and Emerging Diseases*, 67(6), pp. 2336–2344. doi: 10.1111/tbed.13671.
- Lion, T. (2001) 'Current recommendations for positive controls in RT-PCR assays', *Leukemia*, 15(7), pp. 1033–1037. doi: 10.1038/sj.leu.2402133.
- Liu, Y., Meng, H., Shi, L. and Li, L. (2019) 'Sensitive detection of porcine circovirus 3 by droplet digital PCR', *Journal of Veterinary Diagnostic Investigation*, 31(4), pp. 604–607. doi: 10.1177/1040638719847686.
- Matange, K., Tuck, J. M. and Keung, A. J. (2021) 'DNA stability: A central design consideration for DNA data storage systems', *Nature Communications*, 12(1), p. 1358. doi: 10.1038/s41467-021-21587-5.
- Miyazaki, A., Watanabe, S., Ogata, K., Nagatomi, Y., Kokutani, R., Minegishi, Y., Tamehiro, N., Sakai, S., Adachi, R. and Hirao, T. (2019) 'Real-time PCR detection methods for food allergens (wheat, buckwheat, and peanuts) using reference plasmids', *Journal of Agricultural and Food Chemistry*, 67(19), pp. 5680–5686. doi: 10.1021/acs.jafc.9b01234.
- Ouyang, T., Niu, G., Liu, X., Zhang, X., Zhang, Y. and Ren, L. (2019) 'Recent progress on porcine circovirus type 3', *Infection, Genetics and Evolution*, 73, pp. 227–233. doi: 10.1016/j.meegid.2019.05.009.
- Palinski, R., Piñeyro, P., Shang, P., Yuan, F., Guo, R., Fang, Y., Byers, E. and Hause, B. M. (2017) 'A novel porcine circovirus distantly related to known circoviruses is associated with Porcine dermatitis and nephropathy syndrome and

reproductive failure', *Journal of Virology*, 91(1), p. e01873-16. doi: 10.1128/JVI.01879-16.

Phan, T. G., Giannitti, F., Rossow, S., Marthaler, D., Knutson, T. P., Li, L., Deng, X., Resende, T., Vannucci, F. and Delwart, E. (2016) 'Detection of a novel circovirus PCV3 in pigs with cardiac and multi-systemic inflammation', *Virology Journal*, 13(1), p. 184. doi: 10.1186/s12985-016-0642-z.

Saporiti, V., Cruz, T. F., Correa-Fiz, F., Núñez, J. I., Sibila, M. and Segalés, J. (2020a) 'Similar frequency of porcine circovirus 3 (PCV-3) detection in serum samples of pigs affected by digestive or respiratory disorders and age-matched clinically healthy pigs', *Transboundary and Emerging Diseases*, 67(1), pp. 199–205. doi: 10.1111/tbed.13341.

Saporiti, V., Huerta, E., Correa-Fiz, F., Grosse Liesner, B., Duran, O., Segalés, J. and Sibila, M. (2020b) 'Detection and genotyping of porcine circovirus 2 (PCV-2) and detection of porcine circovirus 3 (PCV-3) in sera from fattening pigs of different European countries', *Transboundary and Emerging Diseases*, 67(6), pp. 2521–2531. doi: 10.1111/tbed.13596.

Saporiti, V., Martorell, S., Cruz, T. F., Klaumann, F., Correa-Fiz, F., Balasch, M., Sibila, M. and Segalés, J. (2020c) 'Frequency of detection and phylogenetic analysis of porcine circovirus 3 (PCV-3) in healthy primiparous and multiparous sows and their mummified fetuses and stillborn', *Pathogens*, 9(7), p. 533. doi: 10.3390/pathogens9070533.

Saraiva, G., Vidigal, P., Assao, V., Fajardo, M., Loreto, A., Fietto, J., Bressan, G., Lobato, Z., Almeida, M. and Silva-Júnior, A. (2019) 'Retrospective detection and genetic characterization of porcine circovirus 3 (PCV3) strains identified between 2006 and 2007 in Brazil', *Viruses*, 11(3), p. 201. doi: 10.3390/v11030201.

Serena, M. S., Cappuccio, J. A., Barrales, H., Metz, G. E., Aspitia, C. G., Lozada, I., Perfumo, C. J., Quiroga, M. A., Piñeyro, P. and Echeverría, M. G. (2021) 'First detection and genetic characterization of porcine circovirus type 3 (PCV3) in Argentina and its association with reproductive failure', *Transboundary and Emerging Diseases*, 68(4), pp. 1761–1766. doi: 10.1111/tbed.13893.

Souza, T. C. G. D. de, Gava, D., Schaefer, R., Leme, R. A., Silva Porto, G. da and Alfieri, A. A. (2021) 'Porcine circovirus 3a field strains in free-living wild boars in Paraná State, Brazil', *Animals*, 11(6), p. 1634. doi: 10.3390/ani11061634.

Taverniers, I., Van Bockstaele, E. and De Loose, M. (2004) 'Cloned plasmid DNA fragments as calibrators for controlling GMOs: Different real-time duplex quantitative PCR methods', *Analytical and Bioanalytical Chemistry*, 378(5), pp. 1198–1207. doi: 10.1007/s00216-003-2372-5.

Wang, Y., Feng, Y., Zheng, W., Noll, L., Porter, E., Potter, M., Cino, G., Peddireddi, L., Liu, X., Anderson, G. and Bai, J. (2019) 'A multiplex real-time PCR assay for the detection and differentiation of the newly emerged porcine circovirus type 3 and continuously evolving type 2 strains in the United States', *Journal of Virological Methods*, 269, pp. 7–12. doi: 10.1016/j.jviromet.2019.03.011.

Yao, M., Zhang, X., Gao, Y., Song, S., Xu, D. and Yan, L. (2019) 'Development and application of multiplex PCR method for simultaneous detection of seven viruses in ducks', *BMC Veterinary Research*, 15(1), p. 103. doi: 10.1186/s12917-019-1820-1.

Yuan, L., Liu, Y., Chen, Y., Gu, X., Dong, H., Zhang, S., Han, T., Zhou, Z., Song, X. and Wang, C. (2020) 'Optimized real-time fluorescence PCR assay for the detection of porcine circovirus type 3 (PCV3)', *BMC Veterinary Research*, 16(1), p. 249. doi: 10.1186/s12917-020-02435-y.

Yuzhakov, A. G., Raev, S. A., Alekseev, K. P., Grebennikova, T. V., Verkhovskiy, O. A., Zaberezhny, A. D. and Aliper, T. I. (2018) 'First detection and full genome sequence of porcine circovirus type 3 in Russia', *Virus Genes*, 54(4), pp. 608–611. doi: 10.1007/s11262-018-1582-z.

Zheng, L., Chai, L., Tian, R., Zhao, Y., Chen, H.-Y. and Wang, Z. (2020) 'Simultaneous detection of porcine reproductive and respiratory syndrome virus and porcine circovirus 3 by SYBR Green I-based duplex real-time PCR', *Molecular and Cellular Probes*, 49, p. 101474. doi: 10.1016/j.mcp.2019.101474.